

Risk of Nephrogenic Systemic Fibrosis in Patients With Stage 4 or 5 Chronic Kidney Disease Receiving a Group II Gadolinium-Based Contrast Agent

A Systematic Review and Meta-analysis

Sean A. Woolen, MD, MS; Prasad R. Shankar, MD; Joel J. Gagnier, ND, MSc, PhD; Mark P. MacEachern, MLIS; Lisa Singer, MD, PhD; Matthew S. Davenport, MD

IMPORTANCE Risk of nephrogenic systemic fibrosis (NSF) to individual patients with stage 4 or 5 chronic kidney disease (CKD; defined as estimated glomerular filtration rate of <30 mL/min/1.73 m²) who receive a group II gadolinium-based contrast agent (GBCA) is not well understood or summarized in the literature.

OBJECTIVE To assess the pooled risk of NSF in patients with stage 4 or 5 CKD receiving a group II GBCA.

DATA SOURCES A health sciences informationist searched the Ovid (MEDLINE and MEDLINE Epub Ahead of Print, In-Process & Other Non-Indexed Citation, and Daily and Versions), Embase, Cochrane Central Register of Controlled Trials, Web of Science, and Open Grey databases from inception to January 29, 2019, yielding 2700 citations.

STUDY SELECTION Citations were screened for inclusion in a multistep process. Agreement for final cohort inclusion was determined by 2 blinded screeners using Cohen κ . Inclusion criteria consisted of stage 4 or 5 CKD with or without dialysis, administration of an unconfounded American College of Radiology classification group II GBCA (gadobenate dimeglumine, gadobutrol, gadoterate meglumine, or gadoteridol), and incident NSF as an outcome. Conference abstracts, retracted manuscripts, narrative reviews, editorials, case reports, and manuscripts not reporting total group II GBCA administrations were excluded.

DATA EXTRACTION AND SYNTHESIS Data extraction was performed for all studies by a single investigator, including publication details, study design and time frame, patient characteristics, group II GBCA(s) administered, total exposures for patients with stage 4 or stage 5 CKD, total cases of unconfounded NSF, reason for GBCA administration, follow-up duration, loss to follow-up, basis for NSF screening, and diagnosis.

MAIN OUTCOMES AND MEASURES Pooled incidence of NSF and the associated upper bound of a 2-sided 95% CI (risk estimate) for the pooled data and each of the 4 group II GBCAs.

RESULTS Sixteen unique studies with 4931 patients were included ($\kappa = 0.68$) in this systematic review and meta-analysis. The pooled incidence of NSF was 0 of 4931 (0%; upper bound of 95% CI, 0.07%). The upper bound varied owing to different sample sizes for gadobenate dimeglumine (0 of 3167; upper bound of 95% CI, 0.12%), gadoterate meglumine (0 of 1204; upper bound of 95% CI, 0.31%), gadobutrol (0 of 330; upper bound of 95% CI, 1.11%), and gadoteridol (0 of 230; upper bound of 95% CI, 1.59%).

CONCLUSIONS AND RELEVANCE This study's findings suggest that the risk of NSF from group II GBCA administration in stage 4 or 5 CKD is likely less than 0.07%. The potential diagnostic harms of withholding group II GBCA for indicated examinations may outweigh the risk of NSF in this population.

TRIAL REGISTRATION PROSPERO identifier: [CRD42019123284](https://doi.org/10.1001/jamainternmed.2019.5284)

JAMA Intern Med. doi:10.1001/jamainternmed.2019.5284
Published online December 9, 2019.

[+ Invited Commentary](#)

[+ Supplemental content](#)

Author Affiliations: Department of Radiology, University of Michigan, Ann Arbor (Woolen, Shankar, Davenport); Michigan Radiology Quality Collaborative, University of Michigan, Ann Arbor (Woolen, Shankar, Davenport); Department of Orthopaedic Surgery, University of Michigan, Ann Arbor (Gagnier); Department of Epidemiology, University of Michigan, Ann Arbor (Gagnier); Taubman Health Sciences Library, University of Michigan, Ann Arbor (MacEachern); Dana-Farber Cancer Institute, Department of Radiation Oncology, Brigham and Women's Hospital, Boston, Massachusetts (Singer); Department of Urology, Michigan Medicine, Ann Arbor (Davenport).

Corresponding Author: Prasad R. Shankar, MD, Department of Radiology, University of Michigan, 1500 E Medical Center Dr, Room B2 A209P, Ann Arbor, MI 48109 (pshankar@med.umich.edu).

Nephrogenic systemic fibrosis (NSF) is a rare, potentially fatal condition caused by iatrogenic gadolinium administration in patients with acute kidney injury or stage 4 or 5 chronic kidney disease (CKD), defined as an estimated glomerular filtration rate (eGFR) of less than 30 mL/min/1.73 m².¹⁻⁴ After more than 500 cases of NSF were reported from 1997 to 2007, regulations were adopted to prevent NSF.⁵ In 2007, the US Food and Drug Administration mandated a black box warning advising avoidance of all gadolinium-containing contrast agents (GBCA) in at-risk patients.⁵ The label was updated in 2010 to contain recommendations for health care professionals regarding kidney function screening, use of lower-risk GBCAs, and decreasing GBCA dose.⁶ Such recommendations informed hospital policies and were successful in effectively eliminating the disease.⁷⁻⁹ However, they also resulted in denial or delay of clinically indicated, contrast-enhanced magnetic resonance imaging (MRI) in patients with severe kidney disease, resulting in the undermeasured indirect harms of misdiagnosis and delayed diagnosis.¹⁰ In addition, the guidelines were applied to all GBCAs regardless of gadolinium-chelate lability or association with NSF.^{11,12}

Accumulating literature^{8,13-27} and newer guidelines^{28,29} have recognized that not all GBCAs have the same risk of NSF. The American College of Radiology (ACR) manual on contrast media, version 10.3²⁸ and the European Society of Urogenital Radiology guidelines on contrast agents, version 10.0²⁹ recognize differences in risk of NSF between GBCAs and classify GBCAs into 3 distinct (albeit slightly different) groups.^{28,29} The ACR terms the lowest-risk GBCAs as group II agents (gadobenate dimeglumine, gadoteridol, gadoterate meglumine, and gadobutrol), representing those GBCAs with “very low, if any, risk of NSF development.”^{28(p85)} Both guidelines have been updated recently to indicate that, for the lowest-risk GBCAs, kidney function measurement is not obligatory and that indicated contrast-enhanced MRI with a low-risk GBCA should not be denied on the basis of NSF risk alone.^{28,29}

Unfortunately, the specific risk to individual patients is not well understood or summarized in the literature. Knowledge of this risk is important for counseling and risk-benefit decision-making in individual patients. Establishing these risk estimates may provide an evidence basis for policy makers and physicians who otherwise may hesitate to administer these agents to patients with stage 4 or 5 CKD. The purpose of this systematic review and meta-analysis is to assess the pooled risk of NSF in patients with stage 4 or 5 CKD receiving a group II GBCA.

Methods

This systematic review and meta-analysis was compliant with the Health Insurance Portability and Accountability Act and was exempt based on University of Michigan institutional review board exemption self-regulated status owing to the use of published data with no new study participants. We followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline.³⁰


Key Points

Question What is the risk of nephrogenic systemic fibrosis in patients with stage 4 or 5 chronic kidney disease receiving a group II gadolinium-based contrast agent?

Findings In this systematic review and meta-analysis of 16 unique studies and 4931 patients, the pooled incidence of nephrogenic systemic fibrosis after administration of a group II gadolinium-based contrast agent in patients with stage 4 or 5 chronic kidney disease was 0%; the upper bound of the 95% CI was 0.07%.

Meaning Findings suggest that the risk of nephrogenic systemic fibrosis from group II gadolinium-based contrast agent administration in stage 4 or 5 chronic kidney disease is likely less than 0.07%; potential diagnostic harms of withholding group II gadolinium-based contrast agents for indicated examinations may outweigh the risk of nephrogenic systemic fibrosis in this population.

Figure 1. Study Flow Diagram



Eligibility Criteria

Included studies evaluated human participants with stage 4 or 5 CKD (eGFR, <30 mL/min/1.73 m²) and/or receiving dialysis who underwent administration of a group II GBCA (gadobenate dimeglumine, gadobutrol, gadoterate meglumine, or gadoteridol). The outcome measure required for inclusion was assessment of unconfounded incidence of NSF. All follow-up interval lengths were included. Conference abstracts, retracted manuscripts, narrative reviews, editorials, case reports, and manuscripts not reporting total group II GBCA administrations were excluded (Figure 1).

Data Sources and Searches

Comprehensive searches were performed by an expert health sciences informationist (M.P.M.) from inception to January 29, 2019, in the following databases: Ovid (MEDLINE and MEDLINE Epub Ahead of Print, In-Process & Other Non-Indexed Citation, and Daily and Versions), Embase, Cochrane Central Register of Controlled Trials (Wiley), Web of Science (Clarivate), and Open Grey. Each search consisted of CKD and group II GBCA concept blocks, with combinations of controlled headings (when possible) and title, abstract, and key

Table 1. Characteristics of Included Studies

Source	Country	Study Type	Study Years	Sites	Mean Age, y	No. Female/Total No. (%)
Abujudeh et al, ¹³ 2009	United States	Retrospective cohort	2007-2008	Single	72.6 (SD, 9.6)	152/250 (60.8)
Alhadad et al, ¹⁴ 2012	Sweden	Retrospective cohort	2001-2008	Single	68 (SD, 14)	146/272 (53.7)
Amet et al, ¹⁵ 2014	France	Prospective cohort	2009-2011	Multiple	63 (SD, 14)	Unknown
Bruce et al, ¹⁶ 2016	United States	Retrospective cohort	2006-2014	Single	Unknown	Unknown
Chrysochou et al, ¹⁷ 2010	United Kingdom	Retrospective cohort	1999-2009	Multiple	60.6 (SD, 15.7)	750/2053 (36.5)
Heinz-Peer et al, ¹⁸ 2010	Austria	Retrospective cohort	1997-2007	Single	57.6 (range, 14-91)	79/195 (40.5)
Janus et al, ¹⁹ 2010	France	Retrospective cohort	2005-2006	Multiple	59.9 (range, 18-106)	127/308 (41.2)
Martin et al, ⁸ 2010	United States	Retrospective cohort	2008	Single	51 (range, 17-83)	390/784 (49.7)
Michaely et al, ²⁰ 2017	Multiple	Prospective cohort	2008-2015	Multiple	66.7 (SD, 12.5)	317/908 (34.9)
Nandwana et al, ²¹ 2015	United States	Retrospective cohort	2010-2014	Single	50 (SD, 13)	172/401 (42.9)
Reilly, ²² 2008	United States	Retrospective cohort	2000-2007	Single	61.8 (SD, 9.8)	2/141 (1.4)
Smorodinsky et al, ²³ 2015	United States	Retrospective cohort	2004-2007	Single	53.5 (range, 12-87)	492/1167 (42.2)
Soulez et al, ²⁴ 2015	United States	Prospective cohort	2008-2010	Multiple	63.6 (SD, 13.4) ^a 64.8 (SD, 15.9) ^b	21/45 (46.7) ^a 4/12 (33.3) ^b
Soyer et al, ²⁵ 2017	Multiple	Prospective cohort	2008-2013	Multiple	49.5 (range, 0-98)	18 850/35 499 (53.1)
Tsushima et al, ²⁶ 2018	Japan	Prospective cohort	2015-2017	Multiple	58.1 (SD, 17.4)	1809/3337 (54.2)
Young et al, ²⁷ 2019	United Kingdom	Retrospective cohort	2004-2016	Single	55.6 (SD, 16.1)	8916/15 377 (58.0)

^a Indicates cases of gadobenate dimeglumine administration.

^b Indicates cases of gadoteridol administration.

word terms. No date, language, or other restriction were incorporated into the searches. Duplicate citations were removed in Endnote X8 (Clarivate Analytics). Complete search strategies are available in eMethods 1 in the Supplement.

Study Selection

Studies were screened for inclusion using a multistep process summarized in the study flow diagram (Figure 1). Search results returned 2700 citations, which were screened at the title and abstract level by 2 study team members (1400 by P.R.S. and 1300 by S.A.W.). A sample set of 100 citations was randomly cross-reviewed by the study team member with more years of experience (P.R.S.) to ensure consistency.

After the initial screening, the remaining citations (n = 62) were all reviewed at the manuscript level by 2 blinded study team members (S.A.W. and P.R.S.) (eMethods 2 in the Supplement). Agreement for inclusion was calculated using the Cohen κ with the following scale³¹: 0.01 to 0.20 indicates slight; 0.21 to 0.40, fair; 0.41 to 0.60, moderate; 0.61 to 0.80, substantial; and 0.81 to 0.99, almost perfect. Disagreements were resolved by consensus discussion. A third study team member (M.S.D.) adjudicated when needed.

Data Collection and Data Items

Data extraction was performed for all studies by a single study team member (P.R.S.). Extracted data included publication details, study design and time frame, patient characteristics, group II GBCA(s) administered, total exposures for patients with stage 4 or 5 CKD, total cases of unconfounded NSF, reason for GBCA administration, follow-up duration, loss to follow-up, basis for NSF screening, and diagnosis.

Risk of Bias Analysis

Risk of bias analysis of the included studies was performed by a single study team member (P.R.S.). Criteria used for assessment were based on previously described measures for non-randomized cohort studies.³² Each possible source of bias was assessed as being fulfilled (yes, meaning bias is unlikely to be present), unfulfilled (no, meaning bias is likely to be present), or unknown (meaning information is inadequate or inapplicable to study design). Certain components of the risk of bias assessment were scored as not applicable when inapplicable to study design or results.


Statistical Analysis

The principal summary measure is the pooled incidence of NSF and associated upper bound of the 95% CI (risk estimate) in patients with stage 4 or 5 CKD receiving a group II GBCA. Subanalyses were performed to assess risk estimates on a per-study basis and for each of the 4 individual group II GBCAs. Data analysis was performed with Stata, version 15.2 (StataCorp LLC).

Results

Initial database searches returned 2700 unique citations (Figure 1). After title and abstract review, 62 potential citations remained. After full text review, a final cohort of 16 citations including 4931 patients was available for analysis. Interrater agreement in determining the final study cohort from the 62 screened citations was substantial ($\kappa = 0.68$; 95% CI, 0.49-0.87).

Figure 2. Incidence and Upper Bound of 95% CI of Nephrogenic Systemic Fibrosis (NSF) in Patients With Stage 4 or 5 Chronic Kidney Disease by Study



The 95% CI data are stratified by 16 studies included in the meta-analysis and represent NSF incidence across all studies (0 of 4931 [0%]). Pooled refers to pooled exposures of all studies.

Characteristics of included studies are provided in **Table 1**.^{8,13-27} Studies were published from May 2008 through April 2019. The time frame of investigation across all studies spanned 1997 through 2017. The included studies were a mix of retrospective cohort (11 of 16 [69%]) and prospective cohort (5 of 16 [31%]) designs. Study representation was international, with most of the studies performed in Europe, including 2 multiple-country studies (8 of 16 [50%]) and the United States (7 of 16 [44%]). Multicenter studies constituted 7 of 16 (44%) of the included cohort.

The incidence of NSF in patients with stage 4 or 5 CKD across all 16 studies was 0 of 4931 (0%). The upper bound of the 2-sided 95% CI (1-sided 97.5% CI) for this pooled estimate was 0.07% (**Figure 2**). Study-specific details regarding characteristics of GBCA exposure, number of GBCA exposures, and reference standard for NSF assessment are provided in **Table 2**.

Upper bounds of 95% CIs varied on a study-specific basis (0.26%-52.2%) owing to differences in study-specific eligible sample sizes (**Figure 2**). Follow-up intervals for NSF detection ranged from 3 to 72 months; follow-up interval was unknown in 2 of 16 studies. The reference standard for NSF was most commonly a retrospective medical review (11 of 16 [68.8%]).

The pooled risk of NSF stratified by group II GBCA is provided in **Figure 3**. The greatest safety margin (ie, largest sample size) was for gadobenate dimeglumine (upper bound 95% CI, 0.12% [0 of 3167]). Upper bound 95% CIs for the other group II GBCAs were 1.11% (0 of 330) for gadobutrol, 0.31% (0 of 1204) for gadoterate meglumine, and 1.59% (0 of 230) for gadoteridol.

The risk-of-bias assessment is summarized in the eTable in the **Supplement**. Because the incidence of NSF across all studies was 0%, certain factors in the risk-of-bias assessment related to clustering of outcomes at analysis and adjustment for analysis were not applicable. The most common methodological limitation across the included studies was the unblinding of assessors of NSF to the intervention of GBCA administration (15 of 16 studies [94%]). Strengths of all 16 studies included uniformity in the absence of the outcome (NSF) at the start of investigations, consistency in administration of the intervention (GBCA administration) across all groups, and absence of bias between any potential groups in GBCA administration. A potential for funding bias related to industry support was reported in 7 of 16 studies (44%).

Discussion

Across 16 studies and 4931 administrations, we found the pooled risk of NSF from group II GBCAs in patients with stage 4 or 5 CKD to be 0% (upper bound of 95% CI, 0.07%). This finding indicates the per-patient risk of NSF from group II GBCA administration in stage 4 or 5 CKD is likely less than 0.07%. This risk can be compared with the risk of a severe allergic-like contrast reaction, which has been estimated to be approximately 0.04% for modern low-osmolality iodinated contrast agents³³ and approximately 0.006% to 0.02% for group II GBCAs.³⁴ Despite existing US Food and Drug Administration guidelines indicating that all GBCAs are contraindicated if the eGFR is less than 30 mL/min/1.73 m², these data suggest that group II GBCAs are relatively safe in patients with severe CKD, and their benefits may exceed their risks for indicated examinations. Consistent with our results, recent updates to the ACR,²⁸ European Society of Urogenital Radiology,²⁹ and Canadian Association of Radiologists³⁵ guidelines support use of indicated low-risk GBCAs in this setting.

In comparison with the risk of contrast-induced acute kidney injury, these data indicate that, in patients with stage 4 or 5 CKD who are not receiving dialysis, there is a clearer safety profile for contrast-enhanced MRI using a single-dose group II GBCA than there is for contrast-enhanced computed tomography using a single-dose low-osmolality iodinated contrast agent.^{28,29,36,37} The number needed to harm from low-osmolality iodinated contrast agents (ie, contrast-induced acute kidney injury) has been estimated to be between 1 in 6 and no harm evident (ie, indicating substantial uncertainty) based on recent large, propensity score-adjusted retrospective cohort studies.^{36,37} In both cases (contrast-enhanced computed tomography and contrast-enhanced MRI), the harms of delayed diagnosis and misdiagnosis resulting from the withholding of contrast material in at-risk patients are incompletely measured but likely real.¹⁰ For many disease states, unenhanced imaging has poorer diagnostic accuracy than contrast-enhanced imaging, increasing the risk of diagnostic error and iatrogenic morbidity and mortality.¹⁰

Group II GBCAs include 3 macrocyclic agents with 100% renal excretion (gadoteridol, gadoterate meglumine, and

Table 2. Study Characteristics Related to Group II GBCA Administration and NSF

Source	Total Exposures ^a	GBCA	Reason for GBCA Administration	Follow-up		Standard for NSF	Notes/Comments
				Time	Loss		
Abujudeh et al, ¹³ 2009	6	Gadobenate dimeglumine	Medically necessary	Mean (SD), 3.6 (2.0) mo	None	Medical records and skin/extremity examinations reviewed to identify signs or symptoms of NSF; if no mention, it was assumed the patient did not have NSF	Predominantly patients with stage 3 CKD
Alhadad et al, ¹⁴ 2012	101 (11, 85, and 5)	Multiple (gadobenate dimeglumine, gadoterate meglumine, and gadoteridol)	Routine clinical care	Mean (SD), 46.8 (32.4) mo	None	Electronic medical record for study cohort searched for any sign or symptom of NSF after MRI; diagnosis based on skin biopsy histologic findings	Dermopathologic record review of included cases
Amet et al, ¹⁵ 2014	280 (12, 11, 255, and 2)	Multiple (gadobenate dimeglumine, gadobutrol, gadoterate meglumine, and gadoteridol)	Routine clinical care	At least 4 mo after administration (specific data unavailable)	7.5% Loss (unclear which cohort)	Patient-reported questionnaire of skin findings and evaluation by nephrologist; if suspected, confirmed with 2 dermatologists and 2-site biopsy	Follow-up of patients undergoing dialysis
Bruce et al, ¹⁶ 2016	1423	Gadobenate dimeglumine	Routine clinical practice, protocol to give gadobenate dimeglumine in those with eGFR of <30 mL/min/1.73 m ²	Unknown, follow-up not directly linked to dosing	NA	Institution-wide targeted health care professional survey to assess for known or suspected cases of NSF and annual dermatopathologic review for NSF; diagnosis confirmed based on skin biopsy findings	NSF outcome assessed on a time frame basis, not a per-patient basis
Chrysochou et al, ¹⁷ 2010	483 (445, 13, and 25)	Multiple (gadobenate dimeglumine, gadobutrol, and gadoterate meglumine)	Routine clinical care	Mean (SD), 28.6 (18.2) mo	NA	Medical records of study cohort evaluated for signs of NSF; any skin biopsy records in patients having received gadolinium were evaluated	2278 Patients spanning stages 3-5 CKD, with multiple agents evaluated
Heinz-Peer et al, ¹⁸ 2010	96 (12, 17, 52, and 15)	Multiple (gadobenate dimeglumine, gadobutrol, gadoterate meglumine, and gadoteridol)	Routine clinical care	Unknown	NA	Medical records for study cohort reviewed for documentation of diagnosed NSF or suspected features; histologic findings of any patient with skin biopsy reviewed by dermatologist. Postmortem records reviewed. Diagnosis based on skin biopsy finding or clinical suspicion	Study reports 4 administrations of gadoterate meglumine in confounded cases of NSF
Janus et al, ¹⁹ 2010	135	Gadoterate meglumine	Routine clinical care	Within 4 mo of administration (more precise data unavailable)	NA	Medical records evaluated for cutaneous disorders within 4 mo after MRI; patients routinely evaluated by nephrologist during this interval	3 cases of gadobenate exposure; eGFR data unavailable; no cases of NSF within study cohort
Martin et al, ⁸ 2010	784	Gadobenate dimeglumine	Pretransplant evaluation	Patients included in study if clinical follow-up available 6 mo after last administration; 94%, >10 mo; 6%, 8-10 mo	None	Medical records and dermatopathology records reviewed for all cases in study cohort	All patients had follow-up ≥6 mo
Michaely et al, ²⁰ 2017	284	Gadobutrol	Patients with renal disease requiring contrast-enhanced MRI consented for study inclusion, nonrandomized open label design	Clinical examination at 12 and 24 mo; telephone interviews at 1, 3, 6, and 18 mo after administration	No loss to 2-y record follow-up	Any skin finding of suspected NSF was clinically evaluated; diagnosis based on skin biopsy finding	Analysis stratified by multiple eGFR groups
Nandwana et al, ²¹ 2015	394	Gadobenate dimeglumine	Routine clinical care	>60 d Required for study inclusion; mean, 37.2 mo	None	Electronic medical records reviewed for NSF or NSF-like symptoms	Study details of patients, not exposures; all patients included had follow-up of 60 d
Reilly, ²² 2008	198	Gadoteridol	Routine clinical care	Patients with <14 d excluded; mean (SD), 18.8 (15.6) mo	None	Medical records for study cohort searched for NSF	Veterans Affairs hospital cohort

(continued)


Table 2. Study Characteristics Related to Group II GBCA Administration and NSF (continued)

Source	Total Exposures ^a	GBCA	Reason for GBCA Administration	Follow-up		Standard for NSF	Notes/Comments
				Time	Loss		
Smorodinsky et al, ²³ 2015	40	Gadobenate dimeglumine	Routine clinical care for liver disease	Records were considered sufficient for follow-up if 60 d after administration; mean (range), 49.5 mo (61-3400 d)	19.2% of Cohort	All dermatopathology reports in study patients were reviewed for mention of NSF; manual review of the medical records during study time period also performed	Study evaluating primarily patients in hepatology department
Soulez et al, ²⁴ 2015	50 (40 and 10)	Multiple (gadobenate dimeglumine and gadoteridol)	Routine clinical care	Clinic visits to assess for NSF at 12 and 24 mo after administration; telephone follow-up at 18 mo	5 for Gadobenate dimeglumine and 2 for gadoteridol	Any skin finding of suspected NSF was clinically evaluated by a dermatologist; diagnosis based on skin biopsy finding	Study consisted of 2 parallel studies evaluating gadobenate dimeglumine and gadoteridol
Soyer et al, ²⁵ 2017	65	Gadoterate meglumine	Consecutive eligible patients, routine clinical care	≥3 mo from time of administration to physician follow-up survey; mean, 4.9 mo	7.4%, Unclear which eGFR cohorts	Follow-up questionnaire sent to referring physician to evaluate for signs and symptoms of NSF	Extracted data from much larger cohort study of 35 499 patients
Tsushima et al, ²⁶ 2018	5	Gadobutrol	Noninterventional study of consecutive patients receiving gadobutrol for routine clinical care	3-25 mo for Patients with eGFR <30 mL/min/1.73 m ²	None	Unknown	Extracted data from prospective surveillance study of 3337 patients
Young et al, ²⁷ 2019	587	Gadoterate meglumine	Routine clinical care	Mean (SD), 72 (30) mo	NA	Dermatology records searched during study to identify recorded NSF diagnosis after contrast-enhanced MRI	Only nonconfounded GBCA administrations included in study cohort

Abbreviations: CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; GBCA, gadolinium-based contrast agent; MRI, magnetic resonance imaging; NA, not applicable; NSF, nephrogenic systemic fibrosis.

^a Zero cases of NSF were observed in any of the studies.

Figure 3. Incidence and Upper Bound of 95% CI of Nephrogenic Systemic Fibrosis (NSF) in Patients With Stage 4 or 5 Chronic Kidney Disease by Agent



The 95% CI data are stratified by the 4 gadolinium-based contrast agents included in the study and represent NSF incidence across all studies (0 of 4931 [0%]). Pooled refers to pooled exposures of all agents.

gadobutrol) and 1 linear ionic agent with approximately 95% renal and 5% hepatobiliary excretion (gadobenate dimeglumine).²⁸ Of the 4931 administrations we evaluated, 3167 were exposed to gadobenate dimeglumine. Therefore, most of the safety evidence in this setting is for a specific linear ionic group II GBCA with partial hepatobiliary excretion. Hepatobiliary excretion may offer a protective advantage against NSF by providing an alternative clearance mechanism in patients with severely impaired kidney function.

Previous meta-analyses on GBCA and NSF risk have been heavily weighted by group I GBCAs.^{1,4} Agarwal et al¹ analyzed 7 studies published from 2006 to 2007 with 4276 patients and found an odds ratio of 26.7 (95% CI, 10.3-69.4) for the risk of NSF. Six of the studies evaluated majority or sole group I GBCA exposure, and 1 study had an unknown exposure history. Zhang et al⁴ performed an updated meta-analysis of 11 studies published from 2006 to 2012 with 5405 patients and found an odds ratio of 16.5 (95% CI, 7.5-36.5) for the risk of NSF, suggesting a decline in risk since preventive strategies were introduced. However, because their inclusion criteria required patients diagnosed with NSF, 3 studies of GBCA without evidence of NSF published in 2010, 2013, and 2014 were excluded. Therefore, the existing published meta-analyses^{1,4} are likely not directly relevant to the risk of NSF from group II GBCA.

Strengths and Limitations

Some strengths of our analysis include its focus on a specific and clinically important question, narrow inclusion criteria, a comprehensive search strategy, dual inclusion methods with high interrater agreement, and a low risk of bias for most domains. Common weaknesses included general lack of blinding in the included studies, no universal reference standard for the diagnosis of NSF, and insufficient sample size for specific GBCAs. Most of the studies in our cohort (69%) performed

retrospective evaluations of the medical records to identify potential cases of NSF, raising the possibility that cases could be missed in situations in which individuals were no longer patients in the system where the MRI was performed. However, most studies provided a minimum follow-up interval for study inclusion and provided mean times for record review following GBCA administration (Table 2). In 7 of the studies included in our analysis, development of NSF also was evaluated in patients who received non-group II GBCAs.^{8,14-18,23} In 3 of these studies, NSF was observed.^{8,16,18} These positive controls suggest the lack of NSF detection within our analysis of group II GBCAs was not due solely to methodological biases.

Although our sample size was large (n = 4931), no NSF events occurred. Therefore, the true risk of NSF in this cohort is unknown. The upper bound of the 95% CI was 0.07%, but this result depended on sample size. The absolute risk could be (for example) nonexistent, 1 in a million, or 1 in 2000. With a larger sample, a more precise estimate would be possible. In addition, the analysis reflects studies performed before and after changes to practice guidelines designed to mitigate NSF risk. Therefore, our results are not a pure reflection of either era. There have been single-digit numbers of reports of unconfounded NSF resulting after exposure to a group II GBCA,³⁵ suggesting that the risk of NSF in high-risk patients receiving a group II GBCA is not zero. Larger series are needed to determine what that risk is. Our analysis is unable to determine the risk of sequential group II GBCA exposures or the risk from

group II GBCA administration in the setting of acute kidney injury. The studies we analyzed did not comprehensively or universally address those issues, and this is an area for future investigation. Our analysis was designed to evaluate harms specifically related to development of NSF. It is not a comprehensive assessment of all potential GBCA-related risk (eg, allergic-like reactions, gadolinium retention).

Current ACR guidelines do not require informed consent before group II GBCA administration.²⁸ If a practice wishes to do so, we would suggest the following: “Current evidence does not support withholding group II GBCAs on the basis of NSF risk alone in patients with stage 4 or 5 CKD. Although there is likely a very small risk of developing NSF (likely less than 0.07%) in this population, if the diagnostic question necessitates the use of a GBCA, use of a group II GBCA is recommended.”

Conclusions

The risk of NSF from group II GBCA administration in patients with stage 4 or 5 CKD is likely less than 0.07%. The harms of withholding group II GBCA for indicated examinations may outweigh the risk of NSF in this population. These data support recent updates to ACR and European Society of Urogenital Radiology guidelines^{28,29} liberalizing use of low-risk GBCAs for indicated examinations in this setting.

ARTICLE INFORMATION

Published Online: December 9, 2019.
doi:10.1001/jamainternmed.2019.5284

Author Contributions: Dr Shankar had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Woolen, Shankar, Gagnier, Singer, Davenport.

Acquisition, analysis, or interpretation of data: Woolen, Shankar, Gagnier, MacEachern, Davenport.

Drafting of the manuscript: Woolen, Shankar, Gagnier, MacEachern, Davenport.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Shankar, Gagnier.

Administrative, technical, or material support: Shankar, Gagnier, Singer, Davenport.

Supervision: Shankar, Gagnier, Singer, Davenport.

Conflict of Interest Disclosures: Dr Gagnier reported receiving personal fees from Rubin Anders Scientific and Bartimus Frickleton Robertson Rader PC outside the submitted work. Dr Singer reported receiving grants from Brigham Research Institute outside the submitted work. Dr Davenport reported receiving royalties from Wolters Kluwer (Genitourinary Imaging: A Core Review) and update.com for articles related to iodinated contrast media. No other disclosures were reported.

REFERENCES

1. Agarwal R, Brunelli SM, Williams K, Mitchell MD, Feldman HI, Umscheid CA. Gadolinium-based contrast agents and nephrogenic systemic fibrosis: a systematic review and meta-analysis. *Nephrol Dial*

Transplant. 2009;24(3):856-863. doi:10.1093/ndt/gfn593

2. Gibson SE, Farver CF, Prayson RA. Multiorgan involvement in nephrogenic fibrosing dermopathy: an autopsy case and review of the literature. *Arch Pathol Lab Med.* 2006;130(2):209-212. doi:10.1043/1543-2165(2006)130[209:MIINFD]2.0.CO;2

3. Mendoza FA, Artlett CM, Sandorfi N, Latinis K, Piera-Velazquez S, Jimenez SA. Description of 12 cases of nephrogenic fibrosing dermopathy and review of the literature. *Semin Arthritis Rheum.* 2006;35(4):238-249. doi:10.1016/j.semarthrit.2005.08.002

4. Zhang B, Liang L, Chen W, Liang C, Zhang S. An updated study to determine association between gadolinium-based contrast agents and nephrogenic systemic fibrosis. *PLoS One.* 2015;10(6):e0129720. doi:10.1371/journal.pone.0129720

5. Kitajima K, Maeda T, Watanabe S, Ueno Y, Sugimura K. Recent topics related to nephrogenic systemic fibrosis associated with gadolinium-based contrast agents. *Int J Urol.* 2012;19(9):806-811. doi:10.1111/j.1442-2042.2012.03042.x

6. US Food and Drug Administration. FDA safety communication: new warnings for using gadolinium-based contrast agents in patients with kidney dysfunction. <https://www.fda.gov/Drugs/DrugSafety/ucm223966.htm>. Updated February 6, 2018. Accessed May 9, 2019.

7. Altun E, Martin DR, Wertman R, Lugo-Somolinos A, Fuller ER III, Semelka RC. Nephrogenic systemic fibrosis: change in incidence following a switch in gadolinium agents and adoption of a gadolinium policy—report from two US universities. *Radiology.*

2009;253(3):689-696. doi:10.1148/radiol.2533090649

8. Martin DR, Krishnamoorthy SK, Kalb B, et al. Decreased incidence of NSF in patients on dialysis after changing gadolinium contrast-enhanced MRI protocols. *J Magn Reson Imaging.* 2010;31(2):440-446. doi:10.1002/jmri.22024

9. Wang Y, Alkasab TK, Narin O, et al. Incidence of nephrogenic systemic fibrosis after adoption of restrictive gadolinium-based contrast agent guidelines. *Radiology.* 2011;260(1):105-111. doi:10.1148/radiol.11102340

10. Balogh EPMB, Ball JR; Committee on Diagnostic Error in Health Care; Board on Health Care Services; Institute of Medicine; The National Academies of Sciences, Engineering, and Medicine. *Improving Diagnosis in Health Care.* Washington, DC: National Academies Press; 2015.

11. Abraham JL, Thakral C, Skov L, Rossen K, Marckmann P. Dermal inorganic gadolinium concentrations: evidence for in vivo transmetallation and long-term persistence in nephrogenic systemic fibrosis. *Br J Dermatol.* 2008;158(2):273-280. doi:10.1111/j.1365-2133.2007.08335.x

12. Morcos SK. Extracellular gadolinium contrast agents: differences in stability. *Eur J Radiol.* 2008; 66(2):175-179. doi:10.1016/j.ejrad.2008.01.025

13. Abujudeh HH, Rolls H, Kaewlai R, et al. Retrospective assessment of prevalence of nephrogenic systemic fibrosis (NSF) after implementation of a new guideline for the use of gadobenate dimeglumine as a sole contrast agent for magnetic resonance examination in renally

- impaired patients. *J Magn Reson Imaging*. 2009;30(6):1335-1340. doi:10.1002/jmri.21976
14. Alhadad A, Sterner G, Svensson Å, Alhadad H, Leander P. Incidence of nephrogenic systemic fibrosis at a large university hospital in Sweden. *Scand J Urol Nephrol*. 2012;46(1):48-53. doi:10.3109/00365599.2011.621142
 15. Amet S, Launay-Vacher V, Clément O, et al. Incidence of nephrogenic systemic fibrosis in patients undergoing dialysis after contrast-enhanced magnetic resonance imaging with gadolinium-based contrast agents: the Prospective Fibrose Néphrogénique Systémique study. *Invest Radiol*. 2014;49(2):109-115. doi:10.1097/RLI.0000000000000000
 16. Bruce R, Wentland AL, Haemel AK, et al. Incidence of nephrogenic systemic fibrosis using gadobenate dimeglumine in 1423 patients with renal insufficiency compared with gadodiamide. *Invest Radiol*. 2016;51(11):701-705. doi:10.1097/RLI.0000000000000259
 17. Chrysochou C, Power A, Shurrah AE, et al. Low risk for nephrogenic systemic fibrosis in nondialysis patients who have chronic kidney disease and are investigated with gadolinium-enhanced magnetic resonance imaging. *Clin J Am Soc Nephrol*. 2010;5(3):484-489. doi:10.2215/CJN.06580909
 18. Heinz-Peer G, Neruda A, Watschinger B, et al. Prevalence of NSF following intravenous gadolinium-contrast media administration in dialysis patients with endstage renal disease. *Eur J Radiol*. 2010;76(1):129-134. doi:10.1016/j.ejrad.2009.06.028
 19. Janus N, Launay-Vacher V, Karie S, et al. Prevalence of nephrogenic systemic fibrosis in renal insufficiency patients: results of the FINEST study. *Eur J Radiol*. 2010;73(2):357-359. doi:10.1016/j.ejrad.2008.11.021
 20. Michaely HJ, Aschauer M, Deutschmann H, et al. Gadobutrol in renally impaired patients: results of the GRIP study. *Invest Radiol*. 2017;52(1):55-60. doi:10.1097/RLI.0000000000000307
 21. Nandwana SB, Moreno CC, Osipow MT, Sekhar A, Cox KL. Gadobenate dimeglumine administration and nephrogenic systemic fibrosis: is there a real risk in patients with impaired renal function? *Radiology*. 2015;276(3):741-747. doi:10.1148/radiol.2015142423
 22. Reilly RF. Risk for nephrogenic systemic fibrosis with gadoteridol (ProHance) in patients who are on long-term hemodialysis. *Clin J Am Soc Nephrol*. 2008;3(3):747-751. doi:10.2215/CJN.05721207
 23. Smorodinsky E, Ansdell DS, Foster ZW, et al. Risk of nephrogenic systemic fibrosis is low in patients with chronic liver disease exposed to gadolinium-based contrast agents. *J Magn Reson Imaging*. 2015;41(5):1259-1267. doi:10.1002/jmri.24650
 24. Soulez G, Bloomgarden DC, Rofsky NM, et al. Prospective cohort study of nephrogenic systemic fibrosis in patients with stage 3-5 chronic kidney disease undergoing MRI with injected gadobenate dimeglumine or gadoteridol. *AJR Am J Roentgenol*. 2015;205(3):469-478. doi:10.2214/AJR.14.14268
 25. Soyer P, Dohan A, Patkar D, Gottschalk A. Observational study on the safety profile of gadoterate meglumine in 35,499 patients: the SECURE study. *J Magn Reson Imaging*. 2017;45(4):988-997. doi:10.1002/jmri.25486
 26. Tsushima Y, Awai K, Shinoda G, et al. Post-marketing surveillance of gadobutrol for contrast-enhanced magnetic resonance imaging in Japan. *Jpn J Radiol*. 2018;36(11):676-685. doi:10.1007/s11604-018-0778-4
 27. Young LK, Matthew SZ, Houston JG. Absence of potential gadolinium toxicity symptoms following 22,897 gadoteric acid (Dotarem®) examinations, including 3,209 performed on renally insufficient individuals. *Eur Radiol*. 2019;29(4):1922-1930. doi:10.1007/s00330-018-5737-z
 28. ACR Committee on Drugs and Contrast Media. ACR manual on contrast media, version 10.3. https://www.acr.org/-/media/ACR/Files/Clinical-Resources/Contrast_Media.pdf. Published June 2018. Accessed May 9, 2019.
 29. European Society of Urogenital Radiology. ESUR guidelines on contrast agents, version 10.0. <http://www.esur-cm.org/index.php/a-general-adverse-reactions-2>. Published March 2018. Accessed May 9, 2019.
 30. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-analyses: the PRISMA statement. *Int J Surg*. 2010;8(5):336-341. doi:10.1371/journal.pmed.1000097
 31. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*. 2005;37(5):360-363.
 32. Gagnier JJ, Morgenstern H, Chess L. Interventions designed to prevent anterior cruciate ligament injuries in adolescents and adults: a systematic review and meta-analysis. *Am J Sports Med*. 2013;41(8):1952-1962. doi:10.1177/0363546512458227
 33. Katayama H, Yamaguchi K, Kozuka T, Takashima T, Seez P, Matsuura K. Adverse reactions to ionic and nonionic contrast media: a report from the Japanese Committee on the Safety of Contrast Media. *Radiology*. 1990;175(3):621-628. doi:10.1148/radiology.175.3.2343107
 34. Behzadi AH, Zhao Y, Farooq Z, Prince MR. Immediate allergic reactions to gadolinium-based contrast agents: a systematic review and meta-analysis. *Radiology*. 2018;286(2):471-482. doi:10.1148/radiol.2017162740
 35. Schieda N, Blachman JI, Costa AF, et al. Gadolinium-based contrast agents in kidney disease: a comprehensive review and clinical practice guideline issued by the Canadian Association of Radiologists. *Can J Kidney Health Dis*. 2018;5:2054358118778573. doi:10.1177/2054358118778573
 36. Davenport MS, Cohan RH, Ellis JH. Contrast media controversies in 2015: imaging patients with renal impairment or risk of contrast reaction. *AJR Am J Roentgenol*. 2015;204(6):1174-1181. doi:10.2214/AJR.14.14259
 37. Dekkers IA, van der Molen AJ. Propensity score matching as a substitute for randomized controlled trials on acute kidney injury after contrast media administration: a systematic review. *AJR Am J Roentgenol*. 2018;211(4):822-826. doi:10.2214/AJR.17.19499

Supplementary Online Content

Woolen SA, Shankar PR, Gagnier JJ, MacEachern MP, Singer L, Davenport MS. Risk of nephrogenic systemic fibrosis in patients with stage 4 or 5 chronic kidney disease receiving a group II gadolinium-based contrast agent: a systematic review and meta-analysis. *JAMA Intern Med*. Published online December 9, 2019.
doi:10.1001/jamainternmed.2019.5284

eMethods 1. Complete Search Strategies

eMethods 2. Cohort of 62 Studies Evaluated at Full Manuscript Level by Both Study Team Members

eTable. Risk of Bias Assessment

eReferences.

This supplementary material has been provided by the authors to give readers additional information about their work.

eMethods 1: Complete search strategies

Literature Searches

Ovid MEDLINE; Epub Ahead of Print; In-Process & Other Non-Indexed Citations; Daily and Versions
(1462 results on January 29, 2019)

1. exp renal insufficiency, chronic/ or (renal insufficiency/ and (chronic or end-stage or "stage 4" or "stage 5" or stage iv or stage v).tw,kw) or ((chronic or end-stage or "stage 4" or "stage 5" or stage iv or stage v) adj3 (kidney* or renal)).tw,kw or ((kidney* or renal) adj3 (insufficien* or transplant*)).tw,kw or (impaired adj1 (kidney* or renal)).tw,kw or ((egfr or gfr) adj2 "30").tw,kw or exp renal dialysis/ or dialysis/ or (dialysis or hemodialysis or nsf).ti or nephrogenic fibrosing dermopathy/ or (nephrogenic systemic fibrosis or nephrogenic fibrosing dermopath*).tw,kw
2. ((gadolinium/ or gadolinium dtpa/) and contrast media/) or ((contrast or radiocontrast) adj1 (agent* or dye* or material* or media* or medium*)).ti. or (dotarem or gadavist or gadobenate or gadobutrol or gadoteri* or gadolinium or multihance or prohance).tw,kw
3. (and/1-2) not (exp animals/ not humans/)

Embase.com (1670 results on January 29, 2019)

1. 'chronic kidney failure'/exp OR (((chronic OR 'end stage' OR 'stage iv' OR 'stage v' OR 'stage 4' OR 'stage 5') NEAR/2 (kidney* OR renal)):ti,ab,kw) OR (((kidney* OR renal) NEAR/3 (insufficien* OR transplant*)):ti,ab,kw) OR ((impaired NEAR/1 (kidney* OR renal)):ti,ab,kw) OR (((egfr OR gfr) NEAR/1 '30'):ti,ab,kw) OR dialysis:ti OR hemodialysis:ti OR nsf:ti OR 'nephrogenic systemic fibrosis':ti,ab,kw OR 'nephrogenic fibrosing dermopathies':ti,ab,kw OR 'nephrogenic fibrosing dermopathy':ti,ab,kw
2. 'gadoterate meglumine'/exp OR 'gadobenate dimeglumine'/exp OR 'gadobutrol'/exp OR 'gadoteridol'/exp OR 'gadolinium'/exp OR dotarem:ti,ab,kw OR gadavist:ti,ab,kw OR gadobenate:ti,ab,kw OR gadobutrol:ti,ab,kw OR gadoteri*:ti,ab,kw OR gadolinium:ti,ab,kw OR multihance:ti,ab,kw OR prohance:ti,ab,kw
3. #1 AND #2 NOT ('animal'/exp NOT 'human'/de)

Wiley Cochrane Central Register of Controlled Trials (132 results on January 29, 2019)

1. [mh "renal insufficiency, chronic"] or ([mh "renal insufficiency"] and (chronic or "end-stage" or "stage 4" or "stage 5" or "stage iv" or "stage v"):ti,ab) or ((chronic or "end-stage" or "stage 4" or "stage 5" or "stage iv" or "stage v") near/3 (kidney* or renal)):ti,ab or ((kidney* or renal) near/3 (insufficien* or transplant*)):ti,ab or (impaired near/1 (kidney* or renal)):ti,ab or ((egfr or gfr) near/2 "30"):ti,ab or [mh "renal dialysis"] or [mh dialysis] or (dialysis or hemodialysis or nsf):ti or [mh "nephrogenic fibrosing dermopathy"] or ("nephrogenic systemic fibrosis" or "nephrogenic fibrosing dermopathies" or "nephrogenic fibrosing dermopathy"):ti,ab

2. ([mh gadolinium] or [mh "gadolinium dtpa"]) or ((contrast or radiocontrast) near/1 (agent* or dye* or material* or media* or medium*)):ti or (dotarem or gadavist or gadobenate or gadobutrol or gadoteri* or gadolinium or multihance or prohance):ti,ab

3. #1 and #2

Clarivate Web of Science (836 results on January 29, 2019)

1. TS=(dotarem OR gadavist OR gadobenate OR gadobutrol OR gadoteri* OR multihance OR prohance) OR TS=(gadolinium NEAR/2 contrast*)

2. TS=((chronic OR "end-stage" OR "stage 4" OR "stage 5" OR "stage iv" OR "stage v") NEAR/2 (kidney* or renal)) OR TS=((kidney* OR renal) NEAR/2 (insufficien* OR transplant*)) OR TS=(impaired NEAR/1 (kidney* OR renal)) OR TS=((egfr OR gfr) NEAR/1 "30") OR TI=(dialysis OR hemodialysis OR NSF) OR TS=("nephrogenic systemic fibrosis" OR "nephrogenic fibrosing dermopathies" OR "nephrogenic fibrosing dermopathy")

3. #2 AND #1 NOT TI=(animal* OR bovine OR cattle OR mice OR mouse OR pig OR pigs OR rabbit* OR rat OR rats)

Open Grey (8 results on January 29, 2019)

(dotarem OR gadavist OR gadobenate OR gadobutrol OR gadoteri* OR gadolinium OR multihance OR prohance) AND (chronic OR dialysis OR "end stage" OR nephrogenic OR "stage 4" OR "stage 5" OR "stage iv" OR "stage v")

eMethods 2: Cohort of 62 studies evaluated at full manuscript level by both study team members.

Y. Tsushima, K. Awai, G. Shinoda, H. Miyoshi, M. Chosa, T. Sunaya, J. Endrikat. Post-marketing surveillance of gadobutrol for contrast-enhanced magnetic resonance imaging in Japan. *Japanese Journal of Radiology*. 2018. 36:676-685

D. R. Martin, B. Kalb, A. Mittal, K. Salman, S. Vedantham, P. K. Mittal. Notice of Withdrawal: No Incidence of Nephrogenic Systemic Fibrosis after Gadobenate Dimeglumine Administration in Patients Undergoing Dialysis or Those with Severe Chronic Kidney Disease. *Radiology*. 2018. 286:113-119

J. Endrikat, K. Vogtlaender, S. Dohanish, T. Balzer, J. Breuer. Safety of Gadobutrol: Results From 42 Clinical Phase II to IV Studies and Postmarketing Surveillance After 29 Million Applications. *Investigative Radiology*. 2016. 51:537-43

R. Bruce, A. L. Wentland, A. K. Haemel, R. W. Garrett, D. R. Sadowski, A. Djamali, E. A. Sadowski. Incidence of Nephrogenic Systemic Fibrosis Using Gadobenate Dimeglumine in 1423 Patients With Renal Insufficiency Compared With Gadodiamide. *Investigative Radiology*. 2016. 51:701-705

K. Glutig, R. Bhargava, G. Hahn, W. Hirsch, C. Kunze, H. J. Mentzel, J. F. Schaefer, W. Willinek, P. Palkowitsch, Gardian study group. Safety of gadobutrol in more than 1,000 pediatric patients: subanalysis of the GARDIAN study, a global multicenter prospective non-interventional study. *Pediatric Radiology*. 2016. 46:1317-23

J. R. Costello, B. Kalb, D. R. Martin. Incidence and Risk Factors for Gadolinium-Based Contrast Agent Immediate Reactions. *Topics in Magnetic Resonance Imaging*. 2016. 25:257-263

H. J. Michaely, M. Aschauer, H. Deutschmann, G. Bongartz, M. Gutberlet, R. Woitek, B. Ertl-Wagner, W. Kucharczyk, R. Hammerstingl, F. De Cobelli, M. Rosenberg, T. Balzer, J. Endrikat. Gadobutrol in Renally Impaired Patients: Results of the GRIP Study. *Investigative Radiology*. 2017. 52:55-60

K. M. Shaffer, M. R. Parikh, T. M. Runge, S. D. Perez, S. S. Sakaria, R. M. Subramanian. Renal safety of intravenous gadolinium-enhanced magnetic resonance imaging in patients awaiting liver transplantation. *Liver Transplantation*. 2015. 21:1340-6

B. Zhang, L. Liang, W. Chen, C. Liang, S. Zhang. An Updated Study to Determine Association between Gadolinium-Based Contrast Agents and Nephrogenic Systemic Fibrosis. *PLoS ONE [Electronic Resource]*. 2015. 10:e0129720

L. K. Thomson, P. C. Thomson, D. B. Kingsmore, K. Blessing, C. D. Daly, S. E. Cowper, G. H. Roditi. Diagnosing nephrogenic systemic fibrosis in the post-FDA restriction era. *Journal of Magnetic Resonance Imaging*. 2015. 41:1268-71

E. Smorodinsky, D. S. Ansdell, Z. W. Foster, S. M. Mazhar, I. Cruite, T. Wolfson, S. B. Sugay, G. Iussich, M. Shieh-morteza, Y. Kono, A. Kuo, C. B. Sirlin. Risk of nephrogenic systemic fibrosis is low in patients with

chronic liver disease exposed to gadolinium-based contrast agents. *Journal of Magnetic Resonance Imaging*. 2015. 41:1259-67

G. Soulez, D. C. Bloomgarden, N. M. Rofsky, M. P. Smith, H. H. Abujudeh, D. E. Morgan, R. J. Lichtenstein, M. L. Schiebler, F. J. Wippold, C. Russo, M. J. Kuhn, K. W. Mennitt, J. H. Maki, A. Stolpen, J. Liou, R. C. Semelka, M. A. Kirchin, N. Shen, G. Pirovano, A. Spinazzi. Prospective Cohort Study of Nephrogenic Systemic Fibrosis in Patients With Stage 3-5 Chronic Kidney Disease Undergoing MRI With Injected Gadobenate Dimeglumine or Gadoteridol. *AJR. American Journal of Roentgenology*. 2015. 205:469-78

S. B. Nandwana, C. C. Moreno, M. T. Osipow, A. Sekhar, K. L. Cox. Gadobenate Dimeglumine Administration and Nephrogenic Systemic Fibrosis: Is There a Real Risk in Patients with Impaired Renal Function?. *Radiology*. 2015. 276:741-7

E. Gheuens, R. Daelemans, S. Mesens. Dialysability of gadoteric acid in patients with end-stage renal disease undergoing hemodialysis. *Investigative Radiology*. 2014. 49:505-8

B. J. Edwards, A. E. Laumann, B. Nardone, F. H. Miller, J. Restaino, D. W. Raisch, J. M. McKoy, J. A. Hammel, K. Bhatt, K. Bauer, A. T. Samaras, M. J. Fisher, C. Bull, E. Saddleton, S. M. Belknap, H. S. Thomsen, E. Kanal, S. E. Cowper, A. K. Abu Alfa, D. P. West. Advancing pharmacovigilance through academic-legal collaboration: the case of gadolinium-based contrast agents and nephrogenic systemic fibrosis-a Research on Adverse Drug Events and Reports (RADAR) report. *British Journal of Radiology*. 2014. 87:20140307

B. Nardone, E. Saddleton, A. E. Laumann, B. J. Edwards, D. W. Raisch, J. M. McKoy, S. M. Belknap, C. Bull, A. Haryani, S. E. Cowper, A. K. Abu-Alfa, F. H. Miller, V. Godinez-Puig, V. R. Dharnidharka, D. P. West. Pediatric nephrogenic systemic fibrosis is rarely reported: a RADAR report. *Pediatric Radiology*. 2014. 44:173-80

T. Matsumura, M. Hayakawa, F. Shimada, M. Yabuki, S. Dohanish, P. Palkowitsch, K. Yoshikawa. Safety of gadopentetate dimeglumine after 120 million administrations over 25 years of clinical use. *Magnetic Resonance in Medical Sciences*. 2013. 12:297-304

S. Amet, V. Launay-Vacher, O. Clement, C. Frances, A. Tricotel, B. Stengel, J. Y. Gaurvit, N. Grenier, G. Reinhardt, N. Janus, G. Choukroun, M. Laville, G. Deray. Incidence of nephrogenic systemic fibrosis in patients undergoing dialysis after contrast-enhanced magnetic resonance imaging with gadolinium-based contrast agents: the Prospective Fibrose Nephrogenique Systemique study. *Investigative Radiology*. 2014. 49:109-15

S. G. Schad, P. Heitland, W. N. Kuhn-Velten, G. E. Gross, L. Jonas. Time-dependent decrement of dermal gadolinium deposits and significant improvement of skin symptoms in a patient with nephrogenic systemic fibrosis after temporary renal failure. *Journal of Cutaneous Pathology*. 2013. 40:935-44

B. Winship, S. Ramakrishnan, J. Evans, C. Lurie, D. Martin, T. Pearson, N. Turgeon. A single-center analysis of abdominal imaging in the evaluation of kidney transplant recipients. *Clinical Transplantation*. 2013. 27:701-9

L. P. Le, L. Garibyan, D. Lara, K. E. Finberg, A. J. Iafrate, L. M. Duncan, J. Kay, R. M. Nazarian. Fibrosis-associated single-nucleotide polymorphisms in TGFB1 and CAV1 are not associated with the development of nephrogenic systemic fibrosis. *American Journal of Dermatopathology*. 2013. 35:351-6

G. Deray, O. Rouviere, L. Bacigalupo, B. Maes, T. Hannedouche, F. Vrtovnik, C. Rigother, J. M. Billiow, P. Campioni, J. Ferreiros, D. Devos, D. Alison, F. Glowacki, J. J. Boffa, L. Marti-Bonmati. Safety of meglumine gadoterate (Gd-DOTA)-enhanced MRI compared to unenhanced MRI in patients with chronic kidney disease (RESCUE study). *European Radiology*. 2013. 23:1250-9

S. Becker, S. Walter, O. Witzke, A. Kreuter, A. Kribben, A. Mitchell. Application of gadolinium-based contrast agents and prevalence of nephrogenic systemic fibrosis in a cohort of end-stage renal disease patients on hemodialysis. *Nephron*. 2012. 121:c91-4

C. L. Bennett, K. M. Starko, H. S. Thomsen, S. Cowper, O. Sartor, I. C. Macdougall, Z. P. Qureshi, P. B. Bookstaver, A. D. Miller, L. B. Norris, S. Xirasagar, A. Trenery, I. Lopez, A. Kahn, A. Murday, S. Luminari, D. Cournoyer, F. Locatelli, P. Ray, D. R. Mattison. Linking drugs to obscure illnesses: lessons from pure red cell aplasia, nephrogenic systemic fibrosis, and Reye's syndrome. a report from the Southern Network on Adverse Reactions (SONAR). *Journal of General Internal Medicine*. 2012. 27:1697-703

A. Alhadad, G. Sterner, A. Svensson, H. Alhadad, P. Leander. Incidence of nephrogenic systemic fibrosis at a large university hospital in Sweden. *Scandinavian Journal of Urology & Nephrology*. 2012. 46:48-53

M. Voth, M. Rosenberg, J. Breuer. Safety of gadobutrol, a new generation of contrast agents: experience from clinical trials and postmarketing surveillance. *Investigative Radiology*. 2011. 46:663-71

T. R. Elmholdt, M. Pedersen, B. Jorgensen, K. Sondergaard, J. D. Jensen, M. Ramsing, A. B. Olesen. Nephrogenic systemic fibrosis is found only among gadolinium-exposed patients with renal insufficiency: a case-control study from Denmark. *British Journal of Dermatology*. 2011. 165:828-36

R. A. Rodby. Can gadolinium be given safely to a patient on dialysis?. *Seminars in Dialysis*. 2011. 24:370-1

T. Ishiguchi, S. Takahashi. Safety of gadoterate meglumine (Gd-DOTA) as a contrast agent for magnetic resonance imaging: results of a post-marketing surveillance study in Japan. *Drugs in R & D*. 2010. 10:133-45

D. S. Chow, S. Bahrami, S. S. Raman, S. Rotchel, J. W. Sayre, R. W. Busuttil, D. S. Lu. Risk of nephrogenic systemic fibrosis in liver transplantation patients. *AJR. American Journal of Roentgenology*. 2011. 197:658-62

M. R. Prince, H. Zhang, Z. Zou, R. B. Staron, P. W. Brill. Incidence of immediate gadolinium contrast media reactions. *AJR. American Journal of Roentgenology*. 2011. 196:W138-43

G. Heinz-Peer, A. Neruda, B. Watschinger, A. Vychytil, A. Geusau, M. Haumer, M. Weber. Prevalence of NSF following intravenous gadolinium-contrast media administration in dialysis patients with endstage renal disease. *European Journal of Radiology*. 2010. 76:129-34

C. Chrysochou, A. Power, A. E. Shurrab, S. Husain, S. Moser, J. Lay, A. D. Salama, P. A. Kalra. Low risk for nephrogenic systemic fibrosis in nondialysis patients who have chronic kidney disease and are investigated with gadolinium-enhanced magnetic resonance imaging. *Clinical Journal of The American Society of Nephrology: CJASN*. 2010. 5:484-9

N. Janus, V. Launay-Vacher, S. Karie, O. Clement, E. Ledneva, C. Frances, G. Choukroun, G. Deray. Prevalence of nephrogenic systemic fibrosis in renal insufficiency patients: results of the FINEST study. *European Journal of Radiology*. 2010. 73:357-9

D. R. Martin, S. K. Krishnamoorthy, B. Kalb, K. N. Salman, P. Sharma, J. D. Carew, P. A. Martin, A. B. Chapman, G. L. Ray, C. P. Larsen, T. C. Pearson. Decreased incidence of NSF in patients on dialysis after changing gadolinium contrast-enhanced MRI protocols. *Journal of Magnetic Resonance Imaging*. 2010. 31:440-6

S. M. Mazhar, M. Shiehorteza, C. A. Kohl, M. S. Middleton, C. B. Sirlin. Nephrogenic systemic fibrosis in liver disease: a systematic review. *Journal of Magnetic Resonance Imaging*. 2009. 30:1313-22

Z. Zou, L. Ma, H. Li. Incidence of nephrogenic systemic fibrosis at Chinese PLA General Hospital. *Journal of Magnetic Resonance Imaging*. 2009. 30:1309-12

H. H. Abujudeh, H. Rolls, R. Kaewlai, S. Agarwal, Z. A. Gebreananya, S. Saini, P. W. Schaefer, J. Kay. Retrospective assessment of prevalence of nephrogenic systemic fibrosis (NSF) after implementation of a new guideline for the use of gadobenate dimeglumine as a sole contrast agent for magnetic resonance examination in renally impaired patients. *Journal of Magnetic Resonance Imaging*. 2009. 30:1335-40

H. Wollanka, W. Weidenmaier, C. Giersig. NSF after Gadovist exposure: a case report and hypothesis of NSF development. *Nephrology Dialysis Transplantation*. 2009. 24:3882-4

E. Altun, D. R. Martin, R. Wertman, A. Lugo-Somolinos, E. R. Fuller, R. C. Semelka. Nephrogenic systemic fibrosis: change in incidence following a switch in gadolinium agents and adoption of a gadolinium policy--report from two U.S. universities. *Radiology*. 2009. 253:689-96

R. F. Reilly. Risk for nephrogenic systemic fibrosis with gadoteridol (ProHance) in patients who are on long-term hemodialysis. *Clinical Journal of The American Society of Nephrology: CJASN*. 2008. 3:747-51

B. Tombach, C. Bremer, P. Reimer, F. Matzkies, R. M. Schaefer, W. Ebert, V. Geens, J. Eisele, W. Heindel. Using highly concentrated gadobutrol as an MR contrast agent in patients also requiring hemodialysis: safety and dialysability. *AJR. American Journal of Roentgenology*. 2002. 178:105-9

R. R. Townsend, D. L. Cohen, R. Katholi, S. K. Swan, B. E. Davies, K. Bensef, L. Lambrecht, J. Parker. Safety of intravenous gadolinium (Gd-BOPTA) infusion in patients with renal insufficiency. *American Journal of Kidney Diseases*. 2000. 36:1207-12

S. K. Swan, L. J. Lambrecht, R. Townsend, B. E. Davies, S. McCloud, J. R. Parker, K. Bensef, N. D. LaFrance. Safety and pharmacokinetic profile of gadobenate dimeglumine in subjects with renal impairment. *Investigative Radiology*. 1999. 34:443-8

K. Yoshikawa, A. Davies. Safety of ProHance in special populations. *European Radiology*. 1997. 7 Suppl 5:246-50

L. K. Young, S. Z. Matthew, J. G. Houston. Absence of potential gadolinium toxicity symptoms following 22,897 gadoteric acid (Dotarem) examinations, including 3,209 performed on renally insufficient individuals. *European Radiology*. 2018. 01:01

J. Endrikat, S. Dohanish, N. Schleyer, S. Schwenke, S. Agarwal, T. Balzer. 10 Years of Nephrogenic Systemic Fibrosis: A Comprehensive Analysis of Nephrogenic Systemic Fibrosis Reports Received by a Pharmaceutical Company from 2006 to 2016. *Investigative Radiology*. 2018. 53:541-550

M. R. Prince, J. C. Weinreb. Notice of Withdrawal: MR Imaging and Gadolinium: Reassessing the Risk of Nephrogenic Systemic Fibrosis in Patients with Severe Renal Disease. *Radiology*. 2018. 286:120-121

T. R. Elmholdt, B. Jorgensen, M. Ramsing, M. Pedersen, A. B. Olesen. Two cases of nephrogenic systemic fibrosis after exposure to the macrocyclic compound gadobutrol. *NDT Plus*. 2010. 3:285-287

E. Ozkur, I. Kivanç Altunay, Y. Erdem, G. Özçelik, D. Tuncel. Nephrogenic systemic fibrosis: in a child with primary hyperoxaluria. *Clinical and Experimental Dermatology*. 2019. 44:70-72

D. R. Martin, B. Kalb, A. Mittal, K. Salman, S. Vedantham, P. K. Mittal. No incidence of nephrogenic systemic fibrosis after gadobenate dimeglumine administration in patients undergoing dialysis or those with severe chronic kidney disease. *Radiology*. 2018. 286:113-119

P. Soyer, A. Dohan, D. Patkar, A. Gottschalk. Observational study on the safety profile of gadoterate meglumine in 35,499 patients: The SECURE study. *Journal of Magnetic Resonance Imaging*. 2017. 45:988-997

S. Amet, O. Clément, C. Frances, A. Tricotel, B. Stengel, J. Y. Gauvrit, N. Grenier, G. Reinhardt, N. Janus, L. Rouillon, G. Choukroun, M. Laville, G. Deray, V. Launay-Vacher. French prospective study on nephrogenic systemic fibrosis in hemodialysis patients after contrast media administration: Results of the Pro-FINEST study. *Journal de Pharmacie Clinique*. 2012. 31:149-158

T. Ota, J. Kimura, T. Ishiguchi. Safety and clinical usefulness of gadoteric acid including post-marketing surveillance. *Imaging in Medicine*. 2012. 4:397-409

- R. Renner, R. Treudler, M. Bodendorf, U. Paasch, J. C. Simon. An unusual dermatosclerosis: Nephrogenic systemic fibrosis. *Aktuelle Dermatologie*. 2012. 38:206-208
- M. J. Andrews. Nephrogenic systemic fibrosis: MRI hazards in patients with severe renal impairment. *Journal of the Royal College of Physicians of Edinburgh*. 2008. 38:230-231
- E. Sausseureau, C. Lacroix, A. Cattaneo, L. Mahieu, J. P. Goulle. Erratum to "Hair and fingernail gadolinium ICP-MS contents in an overdose case associated with nephrogenic systemic fibrosis" [Forensic Sci. Int. (in press)] (DOI:10.1016/j.forsciint.2007.06.026). *Forensic Science International*. 2008. 176:91-92
- S. M. Franken, M. Vervloet, M. Starink Th, T. J. Stoof. Gadolinium: A provoking factor of nephrogenic systemic fibrosis?. *Nederlands Tijdschrift voor Dermatologie en Venereologie*. 2007. 17:306-308
- C. M. Kramer, B. J. Schietinger. FDA alert and your practice: MRI contrast agents and nephrogenic systemic fibrosis. *ACC Cardiosource Review Journal*. 2007. 16:48-59
- R. R. Townsend, D. L. Cohen, R. Katholi, S. K. Swan, B. E. Davies, K. Bense, L. Lambrecht, J. Parker. Safety of intravenous Gadolinium (Gd-BOPTA) infusion in patients with renal insufficiency. *Applied Radiology*. 2003. 32:36-41
- D. R. Martin, B. Kalb, A. J. Mittal, K. Salman, S. Vedantham, P. K. Mittal. RETRACTION: No Incidence of Nephrogenic Systemic Fibrosis after Gadobenate Dimeglumine Administration in Patients Undergoing Dialysis or Those with Severe Chronic Kidney Disease (Retraction of Vol 286, Pg 113, 2018). *Radiology*. 2018. 287:363-363
- E. de Kerviler, K. Maravilla, J. F. Meder, O. Naggara, C. Dubourdieu, V. Jullien, P. Desche. Adverse Reactions to Gadoterate Meglumine Review of Over 25 Years of Clinical Use and More Than 50 Million Doses. *Investigative Radiology*. 2016. 51:544-551

eTable 1. Risk of bias assessment

Author *	ROB1	ROB2	ROB3a	ROB3b	ROB4	ROB5	ROB6	ROB7	ROB8	ROB9
Abujudeh HH, et al. (1)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Alhadad A, et al. (2)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Amet S, et al. (3)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Bruce R, et al. (4)	Yes	NA	Yes	Yes	Yes	Yes	Yes	NA	No	Yes
Chrysochou C, et al. (5)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	Yes
Heinz-Peer G, et al. (6)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	Yes
Janus N, et al. (7)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	Yes
Martin DR et al. (8)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	Yes
Michaely HJ, et al. (9)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	No. Funded by Bayer Pharma AG.
Nandwana SB, et al. (10)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Reilly RF, (11)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Smorodinsky E, et al. (12)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	No. At least partial funding from Bracco Group.
Soulez G, et al. (13)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	No. Sponsored by Bracco Diagnostics.
Soyer P, et al. (14)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	Yes
Tsushima Y, et al. (15)	Yes	NA	No	NA	Yes	Yes	Yes	NA	Yes	No. Funded by Bayer Yakuhin, Ltd.
Young LK, et al. (16)	Yes	NA	No	Yes	Yes	Yes	Yes	NA	Yes	Yes

Table legend:

“Yes”: bias is unlikely to be present; “No”: bias is likely to be present; “N/A”: inadequate information or inapplicable to study design

ROB1, Is the outcome absent at the start of the study?

ROB2, Was clustering at the group level accounted for in analyses?

ROB3a., Were the outcome assessors (for the primary outcome) blind to the intervention?

ROB3b., Was the outcome measurement performed in the same manner with similar intensity in the groups being compared?

ROB4, Was a similarly trained individual administering the intervention across groups?

ROB5, Was the outcome measurement performed in the same manner with similar intensity in the groups being compared?

ROB6, Were the groups similar at baseline?

ROB7, Did the authors perform analyses adjusting for known confounders?

ROB8, Were all the withdrawals described?

ROB9, Is the study free of potential funding bias?

eReferences.

1. Abujudeh HH, Rolls H, Kaewlai R, et al. Retrospective assessment of prevalence of nephrogenic systemic fibrosis (NSF) after implementation of a new guideline for the use of gadobenate dimeglumine as a sole contrast agent for magnetic resonance examination in renally impaired patients. *Journal of magnetic resonance imaging : JMRI*. 2009;30(6):1335-1340.
2. Alhadad A, Sterner G, Svensson A, Alhadad H, Leander P. Incidence of nephrogenic systemic fibrosis at a large university hospital in Sweden. *Scandinavian journal of urology and nephrology*. 2012;46(1):48-53.
3. Amet S, Launay-Vacher V, Clement O, et al. Incidence of nephrogenic systemic fibrosis in patients undergoing dialysis after contrast-enhanced magnetic resonance imaging with gadolinium-based contrast agents: the Prospective Fibrose Nephrogenique Systemique study. *Investigative radiology*. 2014;49(2):109-115.
4. Bruce R, Wentland AL, Haemel AK, et al. Incidence of Nephrogenic Systemic Fibrosis Using Gadobenate Dimeglumine in 1423 Patients With Renal Insufficiency Compared With Gadodiamide. *Investigative radiology*. 2016;51(11):701-705.
5. Chrysochou C, Power A, Shurrab AE, et al. Low risk for nephrogenic systemic fibrosis in nondialysis patients who have chronic kidney disease and are investigated with gadolinium-enhanced magnetic resonance imaging. *Clinical journal of the American Society of Nephrology : CJASN*. 2010;5(3):484-489.
6. Heinz-Peer G, Neruda A, Watschinger B, et al. Prevalence of NSF following intravenous gadolinium-contrast media administration in dialysis patients with endstage renal disease. *European journal of radiology*. 2010;76(1):129-134.
7. Janus N, Launay-Vacher V, Karie S, et al. Prevalence of nephrogenic systemic fibrosis in renal insufficiency patients: results of the FINEST study. *European journal of radiology*. 2010;73(2):357-359.
8. Martin DR, Krishnamoorthy SK, Kalb B, et al. Decreased incidence of NSF in patients on dialysis after changing gadolinium contrast-enhanced MRI protocols. *Journal of magnetic resonance imaging : JMRI*. 2010;31(2):440-446.
9. Michaely HJ, Aschauer M, Deutschmann H, et al. Gadobutrol in Renally Impaired Patients: Results of the GRIP Study. *Investigative radiology*. 2017;52(1):55-60.
10. Nandwana SB, Moreno CC, Osipow MT, Sekhar A, Cox KL. Gadobenate Dimeglumine Administration and Nephrogenic Systemic Fibrosis: Is There a Real Risk in Patients with Impaired Renal Function? *Radiology*. 2015;276(3):741-747.
11. Reilly RF. Risk for nephrogenic systemic fibrosis with gadoteridol (ProHance) in patients who are on long-term hemodialysis. *Clinical journal of the American Society of Nephrology : CJASN*. 2008;3(3):747-751.
12. Smorodinsky E, Ansdell DS, Foster ZW, et al. Risk of nephrogenic systemic fibrosis is low in patients with chronic liver disease exposed to gadolinium-based contrast agents. *Journal of magnetic resonance imaging : JMRI*. 2015;41(5):1259-1267.
13. Soulez G, Bloomgarden DC, Rofsky NM, et al. Prospective Cohort Study of Nephrogenic Systemic Fibrosis in Patients With Stage 3-5 Chronic Kidney Disease Undergoing MRI With Injected Gadobenate Dimeglumine or Gadoteridol. *AJR American journal of roentgenology*. 2015;205(3):469-478.

14. Soyer P, Dohan A, Patkar D, Gottschalk A. Observational study on the safety profile of gadoterate meglumine in 35,499 patients: The SECURE study. *Journal of magnetic resonance imaging : JMRI*. 2017;45(4):988-997.
15. Tsushima Y, Awai K, Shinoda G, et al. Post-marketing surveillance of gadobutrol for contrast-enhanced magnetic resonance imaging in Japan. *Japanese journal of radiology*. 2018;36(11):676-685.
16. Young LK, Matthew SZ, Houston JG. Absence of potential gadolinium toxicity symptoms following 22,897 gadoteric acid (Dotarem(R)) examinations, including 3,209 performed on renally insufficient individuals. *European radiology*. 2019;29(4):1922-1930.